

1957

An enzymically catalyzed incorporation of amines into proteins / Nirmal K. Sarkar Donald D. Clarke Heinrich Waelsch Department of Biochemistry, College of Physicians and Surgeons, Columbia University, New York, and the New York Psychiatric Institute, New York, N.Y. (U.S.A.)

Nirmal K. Sarkar

Columbia University. College of Physicians and Surgeons. Department of Biochemistry

Donald Dudley Clarke PhD

Fordham University, clarke@fordham.edu

Heinrich Waelsch

Columbia University. College of Physicians and Surgeons. Department of Biochemistry

Follow this and additional works at: https://fordham.bepress.com/chem_facultypubs

Part of the [Biochemistry Commons](#)

Recommended Citation

Sarkar, Nirmal K.; Clarke, Donald Dudley PhD; and Waelsch, Heinrich, "An enzymically catalyzed incorporation of amines into proteins / Nirmal K. Sarkar Donald D. Clarke Heinrich Waelsch Department of Biochemistry, College of Physicians and Surgeons, Columbia University, New York, and the New York Psychiatric Institute, New York, N.Y. (U.S.A.)" (1957). *Chemistry Faculty Publications*. 40.

https://fordham.bepress.com/chem_facultypubs/40

An enzymically catalyzed incorporation of amines into proteins*

In experiments reported some years ago the enzyme-catalyzed exchange of the amide group of glutamine and asparagine with ^{15}N ammonia was demonstrated but an exchange of the amide groups of protein bound dicarboxylic amino acids could not be explored owing to the unfavorable experimental conditions¹. Recently the problem of the metabolism of protein-amide groups was approached again with the aid of ^{14}C -labeled amines, such as ethanolamine or cadaverine. Soluble protein fractions obtained from sucrose homogenates (0.25 *M*) of the livers of guinea pigs, rats, mice, and rabbits by centrifugation at $100,000 \times g$ were incubated in the presence of Ca^{++} with the ^{14}C -labeled amines. Appreciable radioactivity was recovered in the proteins precipitated and purified by the method of SIEKEVITZ² (Table I). The incorporating system was activated by Ca^{++} but not by Mn^{++} and Mg^{++} or other metal ions—a fact which makes it unlikely that we are dealing with the enzyme system responsible for glutamo or asparto-transferase activities.

TABLE I
INCORPORATION OF ^{14}C -AMINES AND LYSINE INTO SOLUBLE PROTEIN FRACTIONS
OF GUINEA PIG LIVER

Counts/min at infinite thickness

(1) ^{14}C -Amines added	Ethanolamine		Cadaverine		Lysine	
	SP	PP	SP	PP	SP	PP
(2) Protein fraction						
(3) No Ca^{++}	22	530	25	460	23	420
(4) With Ca^{++}	480	5000	690	7270	470	5000
(5) (4) + spermine (4 μmoles)	300	2420	360	3100	330	2500
(6) (4) + hydroxylamine (4 μmoles)	40	120	30	100	50	80
(7) (4) + putrescine (4 μmoles)	270	3060	300	3950	290	3260
(8) (4) + $(\text{NH}_4)^+$ (8 μmoles)	300	3900	340	5000	330	4700

The incubation mixture (1 ml) contained 4 μmoles ^{14}C -amine or ^{14}C -lysine; 20 μmoles CaCl_2 ; 20 μmoles tris(hydroxymethyl)aminomethane, pH 8.0 and 10–12 mg soluble protein (SP) or about 1 mg purified protein (PP). Incubation 1 h at 37° . Incubation was terminated by addition of trichloroacetic acid. Added ^{14}C -lysine and ^{14}C -cadaverine counted 290,000 and 340,000 counts/min at infinite thickness, respectively. ^{14}C -Ethanolamine counted approximately 240,000 counts/min. All protein samples were counted at infinite thickness (11 mg). For counting of experiments with PP, 10–11 mg inert protein was added at the termination of incubation.

BORSOOK *et al.*³ and SCHWEET⁴ have described an enzymically catalyzed incorporation of ^{14}C -lysine into a soluble liver protein fraction. This incorporating system is likewise activated by Ca^{++} and may be identical with the enzyme system active in the incorporation of amines.

Under the conditions of our experiments the soluble liver protein fraction incorporated ^{14}C -lysine as well as ^{14}C -ethanolamine and ^{14}C -cadaverine. The experiments were carried out in tris buffer since phosphate buffer proved to be a strong inhibitor of the incorporation of amines. The pH optimum for the incorporation of the amino acid was found to be 6.6 while that for incorporation of the amines was 8.0 to 8.5. The soluble protein fractions of liver, kidney, and brain of guinea pigs show enzymic activity in incorporating cadaverine in decreasing order. The active protein fraction from guinea pig liver was purified about 10–12 times by ammonium sulfate fractionation and adsorption on calcium phosphate gel and isoelectric precipitation. The purified protein fraction incorporated ^{14}C -cadaverine to an extent of 1 to 2% of the total protein. While nothing definite can be said at present as to the linkage of ^{14}C -cadaverine to the protein, a major portion of the ^{14}C -cadaverine was recovered as mono-dinitrophenyl-cadaverine after hydrolysis of the protein treated with dinitrofluorobenzene.

Inhibition of cadaverine incorporation by other amines (Table I) suggests competition by a variety of amines for the active sites of the protein. Mescaline (4 μmoles) was not inhibitory, a finding of some interest, since it has been reported that it is incorporated into liver proteins⁵.

The occurrence of enzyme systems incorporating biologically occurring amines (and diamino acids) raises the interesting question as to whether or not the enzyme specificity is restricted to aliphatic amines. The findings also open the way to a test whether or not this enzymic reaction,

* This work was supported by grants from the National Institute of Neurological Disease and Blindness (Grant B-226) and from the Ford Foundation.

if occurring *in vivo*, could produce modified body proteins which might interfere with enzymic activities or cell permeability, or might possess antigenic properties.

*Department of Biochemistry, College of Physicians and Surgeons,
Columbia University, New York,
and the New York State Psychiatric Institute, New York, N.Y. (U.S.A.)*

NIRMAL K. SARKAR
DONALD D. CLARKE
HEINRICH WAELSCH

- ¹ H. WAELSCH, P. OWADES, E. BOREK, N. GROSSOWICZ AND M. SCHOU, *Arch. Biochem.*, 27 (1950) 237.
- ² P. SIEKEVITZ, *J. Biol. Chem.*, 195 (1952) 549.
- ³ H. BORSOOK, E. L. DEASY, A. J. HAAGEN-SMIT, G. KEIGHLEY AND P. H. LOWY, *J. Biol. Chem.*, 184 (1950) 529.
- ⁴ R. SCHWEET, *Federation Proc.*, 15 (1956) 350.
- ⁵ W. BLOCK, *Z. physiol. Chem.*, 296 (108) 1954.

Received May 31st, 1957